

Freezing Transitions in Liquids Characterized by a Favoured Local Structure

Pierre Ronceray

Institut de Physique Théorique, CEA Saclay,

F-91191 Gif-sur-Yvette Cedex, France.

Département de Physique, École Normale Supérieure, 75005 Paris

Peter Harrowell

School of Chemistry, University of Sydney, Sydney N.S.W. 2006, Australia

(Dated: December 19, 2012)

Abstract

The ordering transitions of a 2D lattice liquid characterized by a single favoured local structure (FLS) are studied using Monte Carlo simulations. All eight distinct geometries for the FLS are considered and we find a variety of ordering transitions - first order, continuous and multi-step transitions. Using a microcanonical representation of the freezing transition we resolve the dual influence of the local structure on the ordering transition, *i.e.* via the energy of the crystal and the entropy cost of structure in the liquid.

PACS: 61.20.Gy, 64.60.De

I. INTRODUCTION

It is reasonable to presume that among the local structural arrangements that are found to be particularly stable in the liquids will be the ones found in the crystals. It is difficult to see how a liquid could freeze if this was not so. Let us imagine that the crystal is assembled so as to maximise the density of just a single local structure. It would follow that a minimal condition on any Hamiltonian used to model both the liquid and crystal phases is that it must stabilize this particular favoured local structure. The simplest such model would be the one that *only* stabilized this one local structure. This is a simpler picture than is usually invoked for liquid structure. There is, for example, an extended tradition^{1,2} of focussing on non-crystalline favoured local structures in order to explain the kinetic stability of the supercooled liquid with respect to freezing. We note, in passing, that, while reasonable, there is no compelling evidence to suggest that the presence of a competing structure is *necessary* to explain liquid metastability. As we shall see, our simplest case model does not appear to be able to account for the metastability of the supercooled liquid to any significant degree, so providing tacit support for the idea that some sort of competition between multiple stable configurations is required to properly describe the supercooled state. We shall return to this point in the conclusion. In this paper we shall explore the consequences of the minimal model on the liquid and solid states. Our primary interest is to understand how the geometry of the favoured local structure (FLS) determines the nature of the freezing transition.

To address the relationship between the liquid structure and freezing, we shall use a 2D lattice model, the Favoured Local Structure (FLS) model, which allows us to directly select the geometry of a favoured local structure from among the possible geometries³. The benefit of the FLS model is that we do not need to wrestle with the non-trivial task of trying to engineer a desired local geometry through particle interactions. This task has been the subject of a number of recent studies^{4,5}. A specific challenge is that, for small clusters of particles interacting by non-directional short range potentials, the energy proves to be rather insensitive to cluster geometry. This has been addressed by either adding multiple components (up to 8 in an 8 particle cluster⁴) or by exploiting long-range interactions⁵ to select a desired cluster geometry. By assigning energy explicitly based on the cluster geometry itself, as we do in this paper, we effectively set aside the problem of engineering a set of particle interactions to achieve this stability so that we can focus on the consequences

- thermodynamic and kinetic - of the symmetry of a particular favoured local structure.

Previously, we have identified the crystal ground states for each possible choice of FLS³ and demonstrated that the decrease in liquid entropy with decreasing energy is smaller for liquids based on a low symmetry FLS as compared with those based on high symmetry local structures⁶. In this paper, we present an account of the freezing transitions described by the FLS model. In doing so we shall demonstrate how the microcanonical treatment of the phase transitions allows us to clearly see the dual role of the local structure in both establishing the energy of the crystal state and the entropy cost of local ordering in the liquid state.

II. THE FLS MODEL

We consider a 2D triangular lattice of Ising spins (each site has a single degree of freedom named *spin value*, which can take two values called *up* or *down*). Readers should note that this model is identical to one in which two different species A and B are distributed on the lattice. The local environment of a given site is defined as the spin state of its 6 nearest neighbours. In the set of 2^6 possible environments, there are 8 local structures that cannot be inter-converted by rotations or spin inversion. These structures are sketched in the insets on Figure 2. The labeling convention we adopted to design each local structure goes as follow: the first digit is the number of down spin (dark sites on the pictures), and the second one is the length of the longest sequence of up spins in the structure (when there is only 0 or 1 down site, this second digit is unnecessary, and associated structures will be named simply $\{0\}$ and $\{1\}$ respectively). The $\{32\}$ structure has no plane symmetry and is therefore chiral. Here we will consider only the case when the enantiomers are not distinguished. The chiral case $\{32c\}$, where we favour only one enantiomer, involves a number of interesting subtleties that will be addressed in a separate paper⁷.

Among these 8 distinct local structures, we select one that will be designated as the favoured local structure (FLS). Each site whose local environment is the FLS (to a rotation) is assigned an energy of -1 ; all the others have an energy of 0. Note that this energy is independent of the spin on the given site, depending only on the spin arrangement of the 6 neighbouring sites.

We evaluate the equilibrium properties for each FSL model using Monte Carlo sampling. We consider a system in the canonical ensemble, at temperature T (in this work we will

take $k_B = 1$). For high temperatures we have used the standard Metropolis algorithm and, at low temperatures, the equivalent rejection-free method due to Bortz *et al.*⁸ in which sites are organized in terms of the energy change associated with the spin flip.

We have previously³ identified the ground states (all crystalline) for each choice of FLS. To assist the discussion of the various freezing transitions, we present the crystal phases for the different FLS's in Figure 1 along with the energy per site E_o and the number g of distinct realizations of a FLS on the lattice. Note that three FLS's have two distinct crystals with the same energy.

III. LIQUID-CRYSTAL PHASE TRANSITIONS AND THE MICROCANONICAL MAXWELL CONSTRUCTION

The temperature dependence of the average energies on heating and cooling are plotted in Figure 2 for each FLS. We note that the choice of FLS can result in either first order transitions ($\{1\}$, $\{24\}$ and $\{33\}$), continuous transitions ($\{0\}$, $\{22\}$ and $\{31\}$), or, in the case of $\{32\}$ and $\{23\}$, multiple transitions. In the cases where two degenerate polymorphs exist we find that only one of the two is observed to form in the case of $\{23\}$ and $\{24\}$. The observed crystal is labeled a in Figure 1. In the case of $\{31\}$, the polymorphs are spin-inversion symmetric and either can form spontaneously from the liquid.

In addition to the variety of freezing transitions, the equilibrium liquid states exhibit a considerable variation in their heat capacities $C_V = \frac{dE}{dT}$ and the maximum concentration of FLS that the liquid can accumulate before freezing. In a previous paper⁶ we demonstrated that the influence of the choice of FLS on the liquid structure could be attributed to the difference in the entropy cost per FLS. Contributions to this entropy cost per FLS were found to come from both the symmetry of the favoured local structure itself (the higher the symmetry, the greater the entropy cost) and from the entropy of aggregates of FLS's.

Not only does the decrease in symmetry of the FLS allow the liquid to accumulate more local order before freezing, but it also tends to result in large unit cells in the crystal³. This coincidence of effects means that large unit cells in the crystal phase will often be associated with a liquid that can accumulate a significant amount of local structure (i.e. 'pre-order') before freezing and help explain why we see little difference in the rate of crystallization of the different systems, even though the unit cell varies across a factor of 20. To appreciate

the extent of this pre-ordering, consider the case of the $\{32\}$ FLS which freezes into our most complicated crystal structure (see Figure 1. This freezing involves multiple transitions (as we shall describe below) but if we select $T = 0.6$, a temperature that lies just above the first of the ordering transitions, we find that the local ordering has already reached roughly 75% of the final crystal order. In Figure 3 we show an example configuration in the $\{32\}$ liquid at $T = 0.6$ in which the entangled fragments of the crystal order are clearly evident.

Given the established connection between the structure of the FLS and the dependence of entropy $S(E)$ on E in the liquid, it seems natural to consider the freezing transitions from the perspective of the microcanonical ensemble. In this approach, coexistence arises from maximising the entropy at a fixed value of the energy. We can always recover the temperature from the relation

$$\frac{dS}{dE} = \frac{1}{T} = \beta \quad (1)$$

Since both the energy and the entropy are bounded, this approach allows us to embrace the entire temperature range in one plot. In the thermodynamic limit, this construction is completely equivalent to canonical-ensemble constructions.

We can calculate the entropy by integrating $1/T$ over the energy, i.e.

$$S(E) = S(E_i) + \int_{E_i}^E \frac{dE'}{T(E')} \quad (2)$$

where $T(E)$ is the temperature associated to the average energy E in Figure 2. To carry out this calculation we need to know the value of the entropy $S(E_i)$ at the value of the energy E_i from which the integration is carried out. The liquid state is associated to the high temperature limit. At $T \rightarrow \infty$, the entropy per spin is $S_\infty = \ln(2)$ (the factor 2 reflecting the 2 possible states of any individual site) and the energy is the average concentration of FLS's in a random state of the system, i.e.

$$E_\infty = -\frac{g}{2^6} \quad (3)$$

where g is the degeneracy of the FLS and is provided in Figure 1. For the solid entropy, the ground state energy E_0 (also provided in Figure 1) provides the reference energy with an entropy $S(E_0) = 0$. By using Eq. 2 and the respective references states for the liquid and solid, we have calculated the liquid and solid entropies *versus* energy for each choice of

FLS. Any state with an entropy below the convex hull of the $S(E)$ curve corresponds to a metastable state. The common tangent to liquid and crystal entropies therefore corresponds to the thermodynamic freezing transition, and its slope is $1/T_f$. We have constructed this common tangent with the help of fitted curves for the entropy (power law at low temperature, polynomial for the liquid part). This construction is presented on Figure 4 for each choice of FLS.

The transition temperature for the continuous transitions are taken from the peak in the heat capacity. The values of T_f for each of the choices of FLS are presented in Table I along the values of T_f as determined by the Maxwell construction. The common tangent construction is strictly inapplicable to second order transitions. In practical terms, any attempt to identify distinct solid and liquid entropy curves for continuous transitions runs into the problem of sensitivity to the specific choice of energy intervals used to identify the two phases. That said, the Maxwell construction, applied to the continuous transitions, does provide a quite reasonable estimate of the transition temperature.

	Continuous			First order			Multi-step	
FLS	{ 0 }	{ 31 }	{ 22 }	{ 33 }	{ 24 }	{ 1 }	{ 32 }	{ 23 }
T_f	1.43	0.85	0.96	0.69	0.57	0.53	0.34	0.53
							0.50	0.57
							0.56	0.64
$T_{f,\text{maxwell}}$	1.43	0.87	0.97	0.70	0.57	0.54	0.52	0.60

TABLE I: The freezing points of the various FLS systems. In the case of a continuous transition T_f is determined as the position of the maximum of the heat capacity. In the case of a first order transition, T_f is identified as the midpoint between the transition temperatures on heat and cooling. In case of multiple transitions, all the individual transition temperatures are presented. $T_{f,\text{maxwell}}$ was obtained via the common tangent construction as shown in Figure 4.

The microcanonical representation of the freezing transition allows for the most transparent connection to be made between the immediate consequences of the choice of FLS and the transition temperature. To illustrate this connection, we shall consider the following simple treatment of the freezing transition. Let the liquid entropy be given by,

$$S(E) = S_\infty - A(E_\infty - E)^2 \quad (4)$$

As explained in the Appendix, Eq. 4 represents the first term in an expansion of the liquid in terms of $(E_\infty - E)$ and the constant A can be evaluated exactly for this expansion. These calculated values of A are provided in ref.⁶. The second assumption is to neglect the entropy of the crystal, and assume that its entropy at coexistence is zero. This assumption is justified by the observation on Figure 4 that the entropy range for the crystal is indeed very small as compared to the liquid's. A more accurate treatment of the crystal entropy is provided by a single excitation model discussed in the Appendix. With these two assumptions, the common tangent construction is expressed by the relation

$$2A(E_\infty - E) = \frac{S_\infty - A(E_\infty - E)^2}{E - E_o} \quad (5)$$

Solving for E , the value of the liquid energy at coexistence, we have

$$E = E_o + \sqrt{\Delta^2 - S_\infty/A} \quad (6)$$

where $\Delta = E_\infty - E_o$. The theoretical estimate of the transition temperature T_f^{theory} , given by the inverse slope of the common tangent, is

$$T_f^{theory} = \frac{1}{2A(\Delta - \sqrt{\Delta^2 - S_\infty/A})} \quad (7)$$

As is evident from Eqs. 6 and 7, the freezing transition is determined (within this approximate treatment) by just two parameters: the crystal lattice energy, $E_\infty - E_o$, and the liquid entropy factor A . The latter factor contains the entropy cost of structure in the liquid : a large value of A , generally associated with a high symmetry FLS, corresponds to a local structure that incurs a large entropy cost. In Fig. 5 we plot the values of T_f^{theory} calculated using Eq. 7 with the values of the freezing temperature from Table I. Given the simplicity of the treatment, the theoretical estimates of the freezing temperatures are quite reasonable. The notable exception to this success is the FLS {1} where the theory considerably overestimates transition temperature. The {1} liquid actually exhibits a substantial decrease in energy prior to freezing, which is not correctly taken into account by the factor A , as previously noticed⁶, which explains this deviation. While this approximate treatment is too

simple to establish whether a transition is first- or second-order, we note that it does appear to get the magnitude of the transition temperature roughly right, even for continuous transitions.

IV. FREEZING IN MULTIPLE STEPS

The presence of multiple transitions for the $\{23\}$ and $\{32\}$ FLS's is clearly seen when we examine the $E(T)$ curves with an expanded temperature scale about the transition region, as shown in Figure 6. In both cases, the freezing transition is made up of three distinct transitions. This is unusual for crystallization which is typically viewed as the archetype of a highly cooperative transition and hence unlikely to decouple in this way. In this Section we shall try to identify the intermediate phases associated with the step-wise freezing processes.

In the case of the $\{23\}$ FLS, the multiple transition scenario appears to arise from the existence of the two degenerate polymorphs (see Figure 1). If we identify the three transition temperature as T_1 , T_2 and T_3 , as indicated on Figure 6, it appears that the transition on cooling through T_2 corresponds to the formation of a polycrystalline mixture of the two polymorphs. Further cooling sees the formation at T_3 of the single phase of the preferred crystal phase. The nature of the continuous transition at T_1 remains a puzzle.

Turning to the transitions in the $\{32\}$ system, the picture is quite different. Instead of different crystal forms, the sequence of transitions correspond to a step-by-step symmetry breaking. Transition 1 marks the appearance of an orientation, at which the symmetry of the liquid is reduced from “isotropic” to a liquid with 3-fold rotational symmetry. On cooling, transition 2 is characterised by the loss of this rotational symmetry as unique orientation is selected. Finally, transition 3 breaks the reflection symmetry and the translational invariance in one direction, to give the low T crystalline state. Transitions 2 and 3 seem to be first order, though the small energy gap at transition 3 permits oscillation between phases in our finite size systems. This sequence of symmetry breakings is reminiscent of the liquid-nematic-smectic sequence of transitions observed in some molecular liquids.

V. CONCLUSION

In this paper we have examined the variety of freezing transitions exhibited by a model liquid characterised by a favoured local structure. We found examples of ordering transitions that were first order, continuous and multi-stepped. All liquids were found to order with little or, in the case of the continuous transitions, no evidence of a tendency to supercool, in spite of the involvement of some complex crystal phases. The intuitive notion that large unit cell crystals are more 'difficult' to crystallize is challenged here with the demonstration that the very same features that produce such crystals also ensure that a significant amount of the low energy structures will be present in the liquid in advance of the freezing transition.

The microcanonical representation of the freezing transition was presented as the most transparent way of expressing the influence of the local structure on the crystal (through the groundstate energy E_o) and the liquid (via the factor A related to the entropy cost per FLS) on the freezing transition. A simple theoretical treatment that retained only $E_\infty - E_o$ and A was found to provide a reasonable estimate of the transition temperature for most of the liquids considered.

The FLS model involves considerable simplifications. Some, like the restriction to just one FLS, can be removed easily. Others, such as the neglect of the fluctuation of the local structures away from their ideal form cannot easily be included due to the restriction of the lattice. This work poses the question - is there a useful way of expressing, at least approximately, an effective Hamiltonian for interacting particles with continuous degrees of freedom in terms of explicit favoured local structures.

By construction, the crystalline phase of our model is the densest packing of FLS. For all FLS's other than $\{0\}$, these packings don't completely tile the space, a consequence of the fact that these FLS's are frustrated. A large body of work on the notion of geometrical frustration, introduced by Frank in 1952 comment¹, rests on the proposition that the presence of a frustrated FLS can be used to account for the metastability of the supercooled liquid. Based on our results for the FLS model, frustration of an FLS alone is generally insufficient to stabilize the supercooled liquid. The competition between *two* FLS's: one that can produce a low energy crystal while the other FLS, although more stable, can only order in a higher energy structure, looks like the simplest means of generating a glass forming liquid.

While acknowledging the limitations of the FLS model, we believe that it does provide an

excellent tool for building up our intuition on how local structure influences the properties of condensed matter. A report on the extension to 3D is currently in preparation⁹. The role of competing local structures in stabilizing the supercooled liquid and influencing the kinetics of crystallization would involve a straight forward extension of the model presented here. By building the model around the idea of local stable structures, rather than having to discover them as a consequence of particle interactions, we believe that the FLS approach offers a clearer language with which to ask questions about structure in condensed phases - both the influence of local structure and the nature of global ordering.

Acknowledgements

We gratefully acknowledge funding from the École Normale Supérieure and the Australian Research Council.

-
- ¹ F. C. Frank, *Proc. Roy. Soc.* **A215**, 43 (1952).
 - ² G. Tarjus, S. A. Kivelson, Z. Nussinov and P. Viot, *J. Phys.: Cond. Matt.* **17**, R1143 (2005).
 - ³ P. Ronceray and P. Harrowell, *Europhys. Lett.* **96**, 30065 (2011). (2011).
 - ⁴ S. Hormoz and M. P. Brenner, *Proc. Nat. Acad. Sci. USA* **108**, 5193 (2011).
 - ⁵ S. H. Tindemans and B. M. Mulder, *Phys. Rev. E* **82**, 021404 (2010).
 - ⁶ P. Ronceray and P. Harrowell, *J. Chem. Phys.* **136**, 134504 (2012).
 - ⁷ P. Ronceray and P. Harrowell, *in preparation*.
 - ⁸ A. B. Bortz, M. H. Kalos, and J. L. Lebowitz, *J. Comp. Phys.* **17**, 10 (1975).
 - ⁹ P. Ronceray and P. Harrowell, *in preparation*.

APPENDIX

In this Appendix we describe an alternative to the numerical fits used for the common tangent construction, which gives analytic values of the Taylor expansion coefficients around asymptotic high- and low-temperature regimes. We formerly derive the first order term of the expansion⁶. Here we generalize these results to any order using correlation functions that can be computed by exact enumeration. We also present a route to approximate the thermodynamic quantities at low temperature, though only to leading order.

A. The Energy of the Low Temperature Crystal

We shall treat the energy of the crystal phase using a low temperature approximation in which we only include the (uncorrelated) single spin excitations from the groundstate. First, we identify the energy spectrum (n_i, ε_i) of the ground state, as introduced in Table II in the example of system {22}. It classifies the single-spin defects that can occur in the crystal according to the energy of the corresponding state ε_i and the proportion of site that can host such a defect n_i .




State i			
Energy cost ε_i	0 (ground state)	4	6
Concentration n_i	-	3/4	1/4

TABLE II: Sketch of the energy spectrum of system {22} at low temperature. For this system, two kinds of elementary (single spin) defects can occur in the crystal, examples of which are presented schematically (the ground state is also reminded). The energy spectrum of a crystal is determined by the energy costs ε_i of each kind of defect, and the respective concentration of sites that can host such a defect n_i .

In the canonical ensemble at temperature T , each site of type i has a probability:

$$p_i(T) = \frac{e^{-\varepsilon_i/T}}{1 + e^{-\varepsilon_i/T}}$$

to be in its excited state, if we neglect the interaction between such defects. This gives the following approximate expression for the energy :

$$E(T) = E_0 + \sum_i \varepsilon_i n_i \frac{e^{-\varepsilon_i/T}}{1 + e^{-\varepsilon_i/T}} \quad (8)$$

All other thermodynamic quantities derive from $E(T)$. However the computation of $S(T)$, for example, involves complicated analytic forms, and it is much easier to compute it by numerical integration of Eq.8 using Eq. 2, to arbitrary precision.

A quantitative comparison between the results of simulations and this model is presented on Figure 7, for system {22}. We find that the accordance is quantitative up to a temperature around $T = 0.6$. For the other systems, no qualitative difference was observed, though

the quantitative values in the energy spectrum vary. This simple model gives a satisfying description of the crystal state, the physics of which is therefore quite simple. This method gives easily the leading order of the low-temperature expansion. Finding the corrections to this expansion would involve correlation between defects. No difficulty is expected to arise from that, though the computations would be rather tedious.

B. Exact high-temperature expansion of the FLS model at all orders

In this section, we present an adapted version of the classical high-temperature expansions of spin models adapted to the FLS model. It gives the exact values of the coefficients of the Taylor expansion of the thermodynamic functions around $\beta = 1/T = 0$. These coefficients are connected correlation functions at infinite temperature, which may be computed exactly. This reasoning is an extension of our work in⁶, where we presented the route to compute the first non-trivial coefficient.

1. Definitions :

To make the link with other spin models, we first introduce the Hamiltonian (energy of a configuration) :

$$H(\mathcal{C}) = \sum_{i \in \Lambda} \epsilon_i \quad (9)$$

where Λ is the lattice, $\mathcal{C}$ the spin configuration, and the local energy variables are :

$$\epsilon_i = \begin{cases} -1 & \text{if } i \text{ is in the FLS in configuration } \mathcal{C} \\ 0 & \text{otherwise} \end{cases} \quad (10)$$

Each site of the lattice has z neighbours (z is the coordinence of the lattice). We assume invariance under translations, which impose to use periodic boundary conditions if the system is of finite size. For the triangular lattice studied here $z = 6$. Note that the FLS model could be adapted to other regular lattices ; a review of this model on the 3D FCC lattice is in preparation.

The partition function of the system is defined as :

$$\mathcal{Z}(\beta) = \sum_{\mathcal{C}} \exp(-\beta H(\mathcal{C})) \quad (11)$$

where $\beta = 1/T$ is the inverse temperature. All thermodynamic quantities derive from $\mathcal{Z}$ and its derivatives ; for example :

$$F(\beta) = -\frac{1}{\beta} \log \mathcal{Z}(\beta) \quad (12)$$

$$E(\beta) = -\frac{1}{\mathcal{Z}(\beta)} \frac{\partial \mathcal{Z}(\beta)}{\partial \beta} \quad (13)$$

$$S(\beta) = \beta(E - F) \quad (14)$$

are respectively the free energy, the average energy and the entropy of the system.

We define an average $\langle \cdot \rangle_0$ as the uniform average over all configurations, *i.e.* at $\beta = 0$.

Note that for any site i , we have :

$$\langle \epsilon_i \rangle_0 = E_\infty = -\frac{g}{2^z} \quad (15)$$

where g is the *degeneracy* of the FLS and E_∞ is the average energy per site at infinite temperature.

2. Derivation

Now that the notations are set, we are ready to derive an exact Taylor expansion, at any order, of the thermodynamic functions in the FLS model around $\beta = 0$.

Working on the partition function $\mathcal{Z}(\beta)$, a few usual tricks allow us to rewrite:

$$\mathcal{Z}(\beta) = \sum_{\mathcal{C}} \exp(-\beta H(\mathcal{C})) \quad (16)$$

$$= \sum_{\mathcal{C}} \prod_{i \in \Lambda} \exp(\beta \epsilon_i) \quad (17)$$

$$= \sum_{\mathcal{C}} \prod_{i \in \Lambda} (1 + \nu \epsilon_i) \quad (18)$$

with $\nu = e^\beta - 1$. Note that $\nu \approx \beta$ at high temperature. These simplifications use the fact that ϵ_i takes only two values, 0 and 1.

Now we can expand this product exactly :

$$\prod_{i \in \Lambda} (1 + \nu \epsilon_i) = \sum_{A \subseteq \Lambda} \nu^{|A|} \prod_{i \in A} \epsilon_i \quad (19)$$

The meaning of this expansion is the following : A is the subset of right-side factors selected in the expansion of the product, and $|A|$ is the cardinal of A . If we now sum over configurations $\mathcal{C}$:

$$\mathcal{Z}(\beta) = \sum_{\mathcal{C}} \prod_{i \in \Lambda} (1 + \nu \epsilon_i) \quad (20)$$

$$= 2^N \sum_{A \subseteq \Lambda} \nu^{|A|} \langle \prod_{i \in A} \epsilon_i \rangle_0 \quad (21)$$

where N is the total number of sites. We can rearrange this sum according to $|A|$:

$$\mathcal{Z}(\beta) = 2^N \sum_{k \in \mathbf{N}} \sum_{A \subseteq \Lambda, |A|=k} \nu^k \langle \prod_{i \in A} \epsilon_i \rangle_0 \quad (22)$$

So finally we get the high temperature explicit, exact expansion :

$$\boxed{\mathcal{Z}(\beta) = 2^N \sum_{k \in \mathbf{N}} \sum_{i_1 \neq \dots \neq i_k} \nu^k \langle \epsilon_{i_1} \epsilon_{i_2} \dots \epsilon_{i_k} \rangle_0} \quad (23)$$

A convenient rewriting of this equation in terms of the free energy leads us to introduce the connected correlation functions:

$$\boxed{\beta F(\beta) = -\log \mathcal{Z}(\beta) = -N \log(2) - \sum_{k > 0} \sum_{i_1 \neq \dots \neq i_k} \nu^k \langle \epsilon_{i_1} \epsilon_{i_2} \dots \epsilon_{i_k} \rangle_0^C} \quad (24)$$

where :

$$\begin{aligned} \langle \epsilon_i \rangle_0^C &= \langle \epsilon_i \rangle_0 \\ \langle \epsilon_i \epsilon_j \rangle_0^C &= \langle \epsilon_i \epsilon_j \rangle_0 - \langle \epsilon_i \rangle_0 \langle \epsilon_j \rangle_0 \\ \langle \epsilon_i \epsilon_j \epsilon_k \rangle_0^C &= \langle \epsilon_i \epsilon_j \epsilon_k \rangle_0 - \langle \epsilon_i \rangle_0 \langle \epsilon_j \epsilon_k \rangle_0 - \langle \epsilon_j \rangle_0 \langle \epsilon_i \epsilon_k \rangle_0 - \langle \epsilon_k \rangle_0 \langle \epsilon_i \epsilon_j \rangle_0 + 2 \langle \epsilon_i \rangle_0 \langle \epsilon_j \rangle_0 \langle \epsilon_k \rangle_0 \\ &\dots \end{aligned}$$

For the explicit computation, free energy representation is more convenient since connected correlation functions vanish as soon as a site is “far” (not overlapping) from the others. Thus every term in the expansion is extensive in the size of the system. In order to compute explicitly the connected correlation function at order k , we need to enumerate all possible ways to put an FLS in each of the k sites $i_1 \dots i_k$. It will be zero if they can be separated in two non-overlapping set of structures, thus we need only to enumerate a finite number of positions $i_1 \dots i_k$. The quantity $\langle \epsilon_{i_1} \epsilon_{i_2} \dots \epsilon_{i_k} \rangle_0$ can be interpreted simply as the probability that all sites $i_1 \dots i_k$ lie simultaneously in the FLS, in a random spin configuration.

3. Computation of the first terms

The first terms read :

$$\beta F(\beta) = -N \log(2) - \nu \sum_{i \in \Lambda} \langle \epsilon_i \rangle_0^C + \nu^2 \sum_{i \neq j} \langle \epsilon_i \epsilon_j \rangle_0^C + \nu^3 \sum_{i \neq j \neq k} \langle \epsilon_i \epsilon_j \epsilon_k \rangle_0^C + O(\nu^4) \quad (25)$$

The truncature to the order ν^2 would give the Gaussian approximation to the density of states, as in⁶. Now the interest of this new, more formal approach, is to be able to increase the precision and involve higher-order correlation functions.

All that is left is to compute explicitly these connected correlation functions, which need only to test a finite (but possibly big) number of configurations. This is the only model-specific part of this derivation. Using enumerating routines, we could quite easily get the ν^3 term for FCC and triangular lattices. Finally, to obtain an expansion of the free energy in powers of β instead of ν , we can substitute the expansion of ν in terms of β , i.e.

$$\nu = e^\beta - 1 = \sum_{k>0} \frac{\beta^k}{k!} = \beta + \beta^2/2 + \beta^3/6 + \dots \quad (26)$$

into Eq. 25.












FLS	Ground State Crystal		FLS	Ground State Crystal <i>a</i>		Ground State Crystal <i>b</i>	
$\{0\}$ 	$\mathcal{Z} = 1$ $E_0 = -1$ $g = 1$		$\{23\}$ 	$\mathcal{Z} = 9$ $E_0 = -2/3$ $g = 6$		$\mathcal{Z} = 9$ $E_0 = -2/3$	
$\{1\}$ 	$\mathcal{Z} = 7$ $E_0 = -6/7$ $g = 6$		$\{24\}$ 	$\mathcal{Z} = 9$ $E_0 = -2/3$ $g = 6$		$\mathcal{Z} = 3$ $E_0 = -2/3$	
$\{22\}$ 	$\mathcal{Z} = 4$ $E_0 = -3/4$ $g = 3$		$\{31\}$ 	$\mathcal{Z} = 3$ $E_0 = -2/3$ $g = 2$		$\mathcal{Z} = 3$ $E_0 = -2/3$	
$\{32\}$ 	$\mathcal{Z} = 20$ $E_0 = -4/5$ $g = 12$		$\{33\}$ 	$\mathcal{Z} = 6$ $E_0 = -2/3$ $g = 6$			

FIG. 1: The groundstates for the 8 distinct FLS as defined in the text. The geometrical multiplicity g of the FLS, number of sites in the unit cell $\mathcal{Z}$ and the energy per site E_0 in the groundstate are indicated. For the $\{23\}$, $\{24\}$ and $\{31\}$ FLS's we find two degenerate polymorphs.




FIG. 2: The average energy per site as a function of temperature for each FLS. The measurements were made in an hysteresis cycle of temperature -cooling (in blue) then heating (in red). These curves were obtained with a 60×60 lattice and a ‘cooling rate’ of $2 \cdot 10^6$ steps of the rejection-free algorithm per site and per unit of temperature.




FIG. 3: Typical equilibrium spin configuration of the system for FLS $\{32\}$, at temperature $T = 0.6$, just above the first transition. This liquid state is highly structured.




FIG. 4: The entropy $S(E)$ for the crystal (red) and liquid (blue) for each choice of the FLS. The common tangent is indicated as a dashed line in each case.




FIG. 5: A scatter plot of the values of T_f from simulation against the value of T_f^{theory} from Eq. 7. The straight line with slope one indicates the case where $T_f = T_f^{\text{theory}}$.




FIG. 6: Energy versus temperature zoomed curve for multiple transitions systems. The approximate position of the inflexion points are indicated.




FIG. 7: The low temperature model described in equation 8, compared to simulated results for system $\{22\}$. The agreement is excellent at low temperature, as can be seen on the right figure, which is $E(T) - E_0$ plotted in logarithmic scale. This model correctly identifies leading-order behaviour at low temperature.